

Marginal and correlation distribution functions in the squeezed-states representation

Marcelo A. Marchioli*

*Instituto de Física de São Carlos, Universidade de São Paulo,
Caixa Postal 369, 13560-970 São Carlos, SP, Brazil*

Salomon S. Mizrahi† and Victor V. Dodonov‡

*Departamento de Física, Universidade Federal de São Carlos,
Rod. Washington Luiz km 235, 13565-905 São Carlos, SP, Brazil*

(October 20, 1999)

Here we consider the Husimi function P for the squeezed states and calculate the marginal and correlation distribution functions when P is projected onto the photon number states. According to the value of the squeezing parameter one verifies the occurrence of oscillations and beats as already appointed in the literature. We verify that these phenomena are entirely contained in the correlation function. In particular, we show that since the Husimi and its marginal distribution functions satisfy partial differential equations where the squeeze parameter plays the role of time, the solutions (the squeezed functions obtained from “initial” unsqueezed functions) can be expressed by means of kernels responsible for the “propagation” of squeezing. From the calculational point of view, this method presents advantages for calculating the marginal distribution functions (compared to a direct integration over one of the two phase-space variables of P) since one can use the symmetry properties of the differential equations.

PACS number(s): 42.50.Dv, 03.65.Bz, 42.65.Ky

I. INTRODUCTION

In the end of the eighties, the oscillations of the photon distribution function (DF) of high energy squeezed and correlated states were discovered in [1,2]; the authors of [1] studied the oscillatory behavior of the single-mode squeezed-state Husimi function (HF) projected into photon number states, $P_n(p, q; \lambda, \phi) = |\langle n|p, q; \lambda, \phi\rangle|^2$, where p and q are the space variables associated with the two quadratures of a monochromatic electromagnetic (EM) field, λ is the squeezing parameter and ϕ is a rotation angle in phase space. They suggested that for $p = 0$, $q = 7\sqrt{2}$, $\phi = 0$ and a fixed $\lambda = 21$, such oscillatory behavior can be explained in terms of quantum interference effects and were taken as a signature of a nonclassical state. More recently, Gagen [3] generalized this study to incorporate interference structures in the Bohr-Sommerfeld trajectories associated with a superposition of quantum states. On the other hand, Dutta *et al.* [4] verified an additional feature present in $P_n(p, q; \lambda, \phi)$, when plotted as function of n : this distribution exhibits the structure of beats (collapses and revivals) at large values of n (≥ 10) for $\lambda = 201$, $p^2 + q^2 = 98$ and $\phi \approx \pi/2$. Compared to the value λ used in [1], the high value for λ is crucial for the occurrence of beats. These oscillations were attributed to interference effects in phase space [4]; however, since a detailed explanation about the beats has not been presented until the present time, we judge that they deserve a deeper investigation.

Recently, Chountasis and Vourdas [5] showed that the Weyl function is an important tool for quantum interference effects. In particular, they studied the Wigner and Weyl functions for a superposition of m quantum states $|s_i\rangle$, where each function is decomposed into diagonal and nondiagonal terms, where the nondiagonal term is responsible for the interference between the states $|s_i\rangle$. Adopting a different approach and using the formalism developed in [6], and in order to shed more light on the origin of the beats predicted in [4], here we decompose the squeezed state HF into three functions: the two marginal (one for p and the other for q) and the correlation distribution functions (MDF and CDF, respectively). Our results corroborate the phase-space interference concept and complement the graphical treatment proposed by Mandal [7].

*march@ifsc.usp.br

†salomon@power.ufscar.br

‡vdodonov@power.ufscar.br

This paper is organized as follows. In section II we discuss the solution of the differential pseudo-diffusion equation (see reference [6]), and in section III we define the phase space MDF and CDF. The formal and numerical results are given in section IV, where we show that oscillations and beats are present in the CDF function. Section V is dedicated to a summary and conclusions. Two appendices are also presented, containing calculational details. Appendix A contains the main steps to calculate the MDFs by direct integration, and in appendix B we calculate the CDF.

II. THE PSEUDO-DIFFUSION EQUATION AND ITS SOLUTIONS

The mapping of the statistical operator ρ (describing a state of the EM field) on the squeezed-states representation (SSR) permits us to write the HF P as follows [6]:

$$P(p, q; \lambda, \phi) = \text{Tr} [\rho \Pi(p, q; \zeta)] = \text{Tr} [\rho_r \Pi(p_r, q_r; \lambda)] = P_r(p_r, q_r; \lambda) , \quad (1)$$

where

$$\Pi(p, q; \zeta) = |pq; \zeta\rangle\langle pq; \zeta| = \mathbf{R}(\phi/2) \Pi(p_r, q_r; y) \mathbf{R}^\dagger(\phi/2) \quad (2)$$

is a projection operator, $\mathbf{R}(\phi/2) = \exp\left(\frac{i\phi}{2} \mathbf{a}^\dagger \mathbf{a}\right)$ is the rotation operator, $q_r = q \cos(\phi/2) + p \sin(\phi/2)$ and $p_r = p \cos(\phi/2) - q \sin(\phi/2)$ are the rotated phase space variables expressed in terms of the old ones, $\rho_r = \mathbf{R}^\dagger(\phi/2) \rho \mathbf{R}(\phi/2)$ and $\lambda \equiv e^{-2y}$ ($0 < \lambda < \infty$). Now, if one considers the mixed state $\rho = \sum_{n=0}^{\infty} p_n |n\rangle\langle n|$, diagonal in the Fock basis, ρ_r will be invariant under rotations since $\mathbf{R}(\phi/2) |n\rangle = e^{in\phi/2} |n\rangle$. Consequently, the associated HF is given by $P(p, q; \lambda, \phi) = P(p_r, q_r; \lambda)$. This relation is useful in the sense that it is sufficient to consider the calculation of the unrotated distribution $P(p, q; \lambda)$, with variables changed from (p, q) to (p_r, q_r) in the final result, respectively. The number state $|n\rangle\langle n|$ is a typical example where this relation can be used. In [6] we demonstrated that $P(p, q; \lambda, \phi)$ satisfies the partial differential equation

$$\Gamma(p, q; \lambda, \phi) P(p, q; \lambda, \phi) = 0 , \quad (3)$$

where

$$\begin{aligned} \Gamma(p, q; \lambda, \phi) = & \frac{\partial}{\partial \lambda} - \frac{1}{4\lambda^2} \left\{ [\lambda^2 \cos^2(\phi/2) - \sin^2(\phi/2)] \frac{\partial^2}{\partial p^2} + [\lambda^2 \sin^2(\phi/2) - \cos^2(\phi/2)] \frac{\partial^2}{\partial q^2} \right. \\ & \left. - (\lambda^2 + 1) \sin \phi \frac{\partial^2}{\partial q \partial p} \right\} \end{aligned} \quad (4)$$

is a linear differential operator. For $\phi = 0$, equation (3) is similar to the diffusion equation in two dimensions where the parameter λ plays the role of time. In this situation, since the diffusion coefficients have opposite signs, the equation describes a diffusive (infusive) process in the p (q) variable. For this reason, it has been called the *pseudo-diffusion equation* [8,9].

Here we consider the formal solution of equation (3), obtained by the Fourier transform (FT) method, written as an integral equation with the kernel depending on the squeeze and rotation parameters,

$$\begin{aligned} P(p, q; \lambda, \phi) &= \int_{-\infty}^{\infty} \frac{d\xi d\eta}{2\pi} e^{i(\eta p - \xi q)} K(\xi, \eta; \lambda, \phi) \tilde{P}(\xi, \eta) \\ &= \int_{-\infty}^{\infty} \frac{d\xi d\eta}{2\pi} e^{i(\eta p_r - \xi q_r)} K(\xi, \eta; \lambda, 0) \tilde{P}_r(\xi, \eta) \\ &= P_r(p_r, q_r; \lambda) . \end{aligned} \quad (5)$$

The kernel is

$$K(\xi, \eta; \lambda, \phi) = \exp \left(-\frac{\lambda - 1}{4\lambda} \{ [\lambda \sin^2(\phi/2) - \cos^2(\phi/2)] \xi^2 + [\lambda \cos^2(\phi/2) - \sin^2(\phi/2)] \eta^2 + (\lambda + 1) \sin \phi \xi \eta \} \right) \quad (6)$$

and $\tilde{P}(\xi, \eta)$ is the FT of the HF $P(p, q)$ for $\lambda = 1$ and $\phi = 0$. We notice that $K(\xi, \eta; \lambda, \phi)$ is an *unbounded* function, responsible for the squeezing ‘propagation’ of an ‘initial’ function $P(p, q)$ to $P(p, q; \lambda, \phi)$ for any values of λ and ϕ in

their domain. Thus the existence of a ‘propagated’ $P(p, q; \lambda, \phi)$ depends on the functional form of $\tilde{P}(\xi, \eta)$, since the integral in the first line in (5) for $\tilde{P}(\xi, \eta) = \text{constant}$ does not exist. From the pseudo-diffusion equation (3) and the linear differential operator $\mathbf{\Gamma}$, equation (5) shows the following symmetry properties

$$P(p, q; \lambda, \phi) = P(q, -p; \lambda, \phi \pm \pi) = P(q, p; \lambda^{-1}, -\phi) \quad (7)$$

and

$$K(\xi, \eta; \lambda, \phi) = K(\eta, -\xi; \lambda, \phi \pm \pi) = K(\eta, \xi; \lambda^{-1}, -\phi) . \quad (8)$$

Now, our aim is to show that the Glauber-Sudarshan distribution $P^c(p, q; \lambda, \phi)$ and HF $P(p, q; \lambda, \phi)$ are related by

$$P^c(p, q; \lambda, \phi) = \mathbf{\Lambda}(p, q; \lambda, \phi)P(p, q; \lambda, \phi) = P(p, q; -\lambda, \phi) , \quad (9)$$

with

$$\mathbf{\Lambda}(p, q; \lambda, \phi) \equiv \exp \left[-\frac{1}{2} \left(\lambda \frac{\partial^2}{\partial p_r^2} + \lambda^{-1} \frac{\partial^2}{\partial q_r^2} \right) \right] , \quad (10)$$

which can also be written as

$$\mathbf{\Lambda}(p, q; \lambda, \phi) = \exp \left\{ -\frac{1}{2\lambda} \left[[\lambda^2 \cos^2(\phi/2) + \sin^2(\phi/2)] \frac{\partial^2}{\partial p^2} + [\lambda^2 \sin^2(\phi/2) + \cos^2(\phi/2)] \frac{\partial^2}{\partial q^2} - (\lambda^2 - 1) \sin \phi \frac{\partial^2}{\partial p \partial q} \right] \right\} ,$$

since

$$\begin{aligned} \frac{\partial^2}{\partial p_r^2} &= \sin^2(\phi/2) \frac{\partial^2}{\partial q^2} + \cos^2(\phi/2) \frac{\partial^2}{\partial p^2} - \sin \phi \frac{\partial^2}{\partial q \partial p} , \\ \frac{\partial^2}{\partial q_r^2} &= \cos^2(\phi/2) \frac{\partial^2}{\partial q^2} + \sin^2(\phi/2) \frac{\partial^2}{\partial p^2} + \sin \phi \frac{\partial^2}{\partial q \partial p} . \end{aligned}$$

Applying the differential operator $\mathbf{\Lambda}$ on (5), we get

$$\begin{aligned} \mathbf{\Lambda}(p, q; \lambda, \phi)P(p, q; \lambda, \phi) &= \int_{-\infty}^{\infty} \frac{d\xi d\eta}{2\pi} \left[\mathbf{\Lambda}(p, q; \lambda, \phi) e^{i(\eta p_r - \xi q_r)} \right] K(\xi, \eta; \lambda, 0) \tilde{P}_r(\xi, \eta) \\ &= \int_{-\infty}^{\infty} \frac{d\xi d\eta}{2\pi} e^{i(\eta p_r - \xi q_r)} \underbrace{e^{\frac{1}{2}(\lambda^{-1}\xi^2 + \lambda\eta^2)} K(\xi, \eta; \lambda, 0)}_{K(\xi, \eta; -\lambda, 0)} \tilde{P}_r(\xi, \eta) \\ &= \int_{-\infty}^{\infty} \frac{d\xi d\eta}{2\pi} e^{i(\eta p_r - \eta q_r)} K(\xi, \eta; -\lambda, 0) \tilde{P}_r(\xi, \eta) \\ &= P(p, q; -\lambda, \phi) . \end{aligned} \quad (11)$$

The second equality is obtained using the following relation

$$\begin{aligned} \mathbf{\Lambda}(p, q; \lambda, \phi) e^{i(\eta p_r - \xi q_r)} &= \sum_{k=0}^{\infty} \frac{(-1)^k}{2^k k!} \left(\lambda^{-1} \frac{\partial^2}{\partial q_r^2} + \lambda \frac{\partial^2}{\partial p_r^2} \right)^k e^{i(\eta p_r - \xi q_r)} \\ &= \sum_{k=0}^{\infty} \frac{1}{k!} \left[\frac{1}{2} (\lambda^{-1} \xi^2 + \lambda \eta^2) \right]^k e^{i(\eta p_r - \xi q_r)} \\ &= e^{\frac{1}{2}(\lambda^{-1} \xi^2 + \lambda \eta^2)} e^{i(\eta p_r - \xi q_r)} \end{aligned}$$

and, from the definition of the kernel $K(\xi, \eta; \lambda, 0)$, we conclude that

$$e^{\frac{1}{2}(\lambda^{-1} \xi^2 + \lambda \eta^2)} K(\xi, \eta; \lambda, 0) = e^{\frac{1}{4}(\lambda+1)(\eta^2 + \lambda^{-1} \xi^2)} = K(\xi, \eta; -\lambda, 0) .$$

Thus, the DF $P^c(p, q; \lambda, \phi)$ is directly obtained by changing the signal of the squeezing parameter $\lambda \rightarrow -\lambda$ in the HF. Consequently, this result shows that $P_n^c(p, q; \lambda, \phi)$ does not exist as a bounded function for the number states, however it exists as an ultradistribution. Equation (9) is a generalization of previous results obtained in [9,10] for $\phi = 0$.

III. MARGINAL AND CORRELATION DISTRIBUTION FUNCTIONS

The HF $P(p, q; \lambda, \phi)$ can be written as a sum of two terms: the first is the product of the two MDF, in q and p , and describes the noncorrelated part; the second term is the CDF and contains the phase-space correlations [6]. So, the HF (1) can be written as

$$P(p, q; \lambda, \phi) = R(p; \lambda, \phi)Q(q; \lambda, \phi) + C(p, q; \lambda, \phi) , \quad (12)$$

where

$$Q(q; \lambda, \phi) = \int_{-\infty}^{\infty} \frac{dp}{\sqrt{2\pi}} P(p, q; \lambda, \phi) , \quad (13)$$

and

$$R(p; \lambda, \phi) = \int_{-\infty}^{\infty} \frac{dq}{\sqrt{2\pi}} P(p, q; \lambda, \phi) \quad (14)$$

are the MDFs, and $C(p, q; \lambda, \phi)$ is the CDF. However, the calculation of (13) and (14) by direct integration displays difficulties when $\phi \neq 0$ (see appendix A). Thus, the aim of this section is to show that expressions for the MDFs can be obtained in a much simpler way by using the formalism of section II.

Substituting the right hand side (RHS) of the first line of equation (5) into the integrands of equations (13) and (14), and then carrying out the integrations we get

$$Q(q; \lambda, \phi) = \int_{-\infty}^{\infty} \frac{d\xi}{\sqrt{2\pi}} e^{-i\xi q} k_Q(\xi; \lambda, \phi) \tilde{Q}(\xi) \quad (15)$$

and

$$R(p; \lambda, \phi) = \int_{-\infty}^{\infty} \frac{d\eta}{\sqrt{2\pi}} e^{i\eta p} k_R(\eta; \lambda, \phi) \tilde{R}(\eta) , \quad (16)$$

where

$$k_Q(\xi; \lambda, \phi) = \exp \left\{ -\frac{\lambda-1}{4\lambda} [\lambda \sin^2(\phi/2) - \cos^2(\phi/2)] \xi^2 \right\} \quad (17)$$

and

$$k_R(\eta; \lambda, \phi) = \exp \left\{ -\frac{\lambda-1}{4\lambda} [\lambda \cos^2(\phi/2) - \sin^2(\phi/2)] \eta^2 \right\} \quad (18)$$

are the reduced kernels responsible for the ‘propagation’ of the squeezing. The functions $\tilde{Q}(\xi)$ and $\tilde{R}(\eta)$ are the respective FT of the Husimi functions $Q(q)$ and $R(p)$ for $\lambda = 1$ (absence of squeezing). Furthermore, the equations (15) and (16) are solutions of the partial differential equations

$$\left[\frac{\partial}{\partial \lambda} - \frac{\lambda^2 \sin^2(\phi/2) - \cos^2(\phi/2)}{4\lambda^2} \frac{\partial^2}{\partial q^2} \right] Q(q; \lambda, \phi) = 0 , \quad (19)$$

$$\left[\frac{\partial}{\partial \lambda} - \frac{\lambda^2 \cos^2(\phi/2) - \sin^2(\phi/2)}{4\lambda^2} \frac{\partial^2}{\partial p^2} \right] R(p; \lambda, \phi) = 0 . \quad (20)$$

In analogy to (8), the reduced kernels k_Q and k_R have the symmetry properties

$$k_{Q(R)}(x; \lambda, \phi) = k_{R(Q)}(x; \lambda^{-1}, \phi) = k_{R(Q)}(x; \lambda, \pi \pm \phi) , \quad (21)$$

which reflect directly into the MDFs,

$$\begin{aligned} Q(q; \lambda, \phi) &= R(q; \lambda^{-1}, \phi) = R(q; \lambda, \pi \pm \phi) , \\ R(p; \lambda, \phi) &= Q(p; \lambda^{-1}, \phi) = Q(p; \lambda, \pi \pm \phi) . \end{aligned} \quad (22)$$

Consequently, the calculation of $Q(q; \lambda, \phi)$ is sufficient for determining the function $R(p; \lambda, \phi)$, and vice-versa.

Now we analyze the structure of the kernel (6), which can be factorized as

$$K(\xi, \eta; \lambda, \phi) = k_Q(\xi; \lambda, \phi) k_R(\eta; \lambda, \phi) k_C(\xi, \eta; \lambda, \phi) , \quad (23)$$

where the first two factors on the RHS (reduced kernels) ‘propagate’ the initial HF in an independent way, *i.e.*, if in the ‘initial’ ($\lambda = 1$) HF the phase-space variables are not correlated, they will remain as such for any other value of λ . The factor

$$k_C(\xi, \eta; \lambda, \phi) = \exp \left[- \left(\frac{\lambda^2 - 1}{4\lambda} \sin \phi \right) \xi \eta \right] \quad (24)$$

introduces additional (or new) correlations into an ‘initial’ HF when $\phi \neq n\pi$, with $n \in \mathbb{N}$. Otherwise, we obtain $k_C = 1$ and $K(\xi, \eta; \lambda, n\pi) = k_Q(\xi; \lambda, n\pi) k_R(\eta; \lambda, n\pi)$, one reduced kernel for each variable.

As a consequence of the factorization (23) it is interesting to rewrite the CDF $C(p, q; \lambda, \phi)$ as a sum of two terms,

$$C(p, q; \lambda, \phi) = C^{(1)}(p, q; \lambda, \phi) + C^{(2)}(p, q; \lambda, \phi) , \quad (25)$$

defined as

$$C^{(1)}(p, q; \lambda, \phi) = \int_{-\infty}^{\infty} \frac{d\xi d\eta}{2\pi} e^{i(\eta p - \xi q)} K(\xi, \eta; \lambda, \phi) \tilde{C}(\xi, \eta) \quad (26)$$

and

$$C^{(2)}(p, q; \lambda, \phi) = \int_{-\infty}^{\infty} \frac{d\xi d\eta}{2\pi} e^{i(\eta p - \xi q)} k_Q(\xi; \lambda, \phi) k_R(\eta; \lambda, \phi) [k_C(\xi, \eta; \lambda, \phi) - 1] \tilde{R}(\eta) \tilde{Q}(\xi) . \quad (27)$$

Here we assume that the HF contains ‘initial’ correlations [see equation (12)] with its FT being $\tilde{P}(\xi, \eta) = \tilde{R}(\eta) \tilde{Q}(\xi) + \tilde{C}(\xi, \eta)$, and that ‘propagation’ of correlations originates from two sources. The first, the RHS of equation (26), is responsible for the ‘propagation’ of squeezing into the ‘initial’ correlations $\tilde{C}(\xi, \eta)$. In the second, equation (27), ‘propagation’ occurs only for $\phi \neq n\pi$ when additional correlations are created into the ‘initial’ FT of the uncorrelated part of the HF, $\tilde{R}(\eta) \tilde{Q}(\xi)$. In appendix B, the correlations $C^{(1)}$ and $C^{(2)}$ are obtained for the Fock states in the SSR.

IV. FOCK STATES IN THE SQUEEZED STATES REPRESENTATION

The density operator $\rho_n = |n\rangle\langle n|$ mapped in the coherent states representation yields a Poisson distribution [11]

$$P_n(p, q) = |\langle pq|n\rangle|^2 = \frac{1}{n!} \left(\frac{p^2 + q^2}{2} \right)^n \exp \left(-\frac{p^2 + q^2}{2} \right) , \quad (28)$$

with $n = 0, 1, 2, \dots$. So, the respective Fourier transform

$$\tilde{P}_n(\xi, \eta) = \frac{(-1)^n}{2^{2n} n!} \sum_{k=0}^n \frac{n!}{k!(n-k)!} \mathcal{H}_{2k} \left(\frac{\xi}{\sqrt{2}} \right) \mathcal{H}_{2(n-k)} \left(\frac{\eta}{\sqrt{2}} \right) \exp \left(-\frac{\xi^2 + \eta^2}{2} \right) , \quad (29)$$

where $\mathcal{H}_m(x)$ is the Hermite polynomial, represents an initial step for calculating the Husimi function $P_n(p, q; \lambda, \phi)$ in the squeezed states representation. In fact, substituting equation (29) into (5) and evaluating the integrations over ξ and η [12, §7.374-8], we get

$$P_n(p, q; \lambda, \phi) = \frac{2\sqrt{\lambda}}{\lambda + 1} \left(\frac{\lambda - 1}{\lambda + 1} \right)^n \frac{1}{2^n n!} \left| \mathcal{H}_n \left(\frac{\lambda q_r + i p_r}{\sqrt{\lambda^2 - 1}} \right) \right|^2 \exp \left(-\frac{\lambda q_r^2 + p_r^2}{\lambda + 1} \right) , \quad (30)$$

where q_r and p_r are the rotated variables defined in section II. This expression was initially obtained in [13], and later used by Schleich *et al.* [1] in the oscillatory behavior study of the distribution $P_n(p, q; \lambda, 0)$. Now, using the mathematical relation [14]

$$|\mathcal{H}_n(z)|^2 = 2^n n! \sum_{k=0}^n (-1)^k \mathcal{L}_k^{(-1/2)}(2x^2) \mathcal{L}_{n-k}^{(-1/2)}(-2y^2) \quad (z = x + iy) ,$$

in which $\mathcal{L}_m^{(\alpha)}(x)$ is the associated Laguerre polynomial, equation (30) can be written in an equivalent form,

$$P_n(p, q; \lambda, \phi) = \frac{2\sqrt{\lambda}}{\lambda+1} \left(\frac{\lambda-1}{\lambda+1} \right)^n \sum_{k=0}^n (-1)^k \mathcal{L}_k^{(-1/2)} \left(\frac{2\lambda^2 q_r^2}{\lambda^2 - 1} \right) \mathcal{L}_{n-k}^{(-1/2)} \left(-\frac{2p_r^2}{\lambda^2 - 1} \right) \exp \left(-\frac{\lambda q_r^2 + p_r^2}{\lambda+1} \right). \quad (31)$$

The Husimi function $Q_n(q)$ is obtained with the help of equation (28), *i.e.*,

$$Q_n(q) = \int_{-\infty}^{\infty} \frac{dp}{\sqrt{2\pi}} P_n(p, q) = \exp \left(-\frac{q^2}{2} \right) \sum_{k=0}^n \mathcal{L}_{n-k}^{(-1/2)}(0) \frac{q^{2k}}{2^k k!}, \quad (32)$$

whose Fourier transform is given by

$$\tilde{Q}_n(\xi) = \sum_{k=0}^n \mathcal{L}_{n-k}^{(-1/2)}(0) \mathcal{L}_k^{(-1/2)} \left(\frac{\xi^2}{2} \right) \exp \left(-\frac{\xi^2}{2} \right). \quad (33)$$

Substituting this result into equation (15) and doing the integration with respect to ξ , we get

$$Q_n(q; \lambda, \phi) = \sqrt{\frac{2\lambda}{(\lambda+1)[\cos^2(\phi/2) + \lambda \sin^2(\phi/2)]}} \sum_{k=0}^n (-1)^k \mathcal{L}_{n-k}^{(-1/2)}(0) \left[\frac{\lambda-1}{\lambda+1} \frac{\cos^2(\phi/2) - \lambda \sin^2(\phi/2)}{\cos^2(\phi/2) + \lambda \sin^2(\phi/2)} \right]^k \times \mathcal{L}_k^{(-1/2)} \left[\frac{2\lambda^2 q^2}{(\lambda^2 - 1)[\cos^4(\phi/2) - \lambda^2 \sin^4(\phi/2)]} \right] \exp \left[-\frac{\lambda q^2}{(\lambda+1)[\cos^2(\phi/2) + \lambda \sin^2(\phi/2)]} \right]. \quad (34)$$

In order to obtain the MDF $R_n(p; \lambda, \phi)$, we only need the symmetry properties (22),

$$R_n(p; \lambda, \phi) = \sqrt{\frac{2\lambda}{(\lambda+1)[\lambda \cos^2(\phi/2) + \sin^2(\phi/2)]}} \sum_{k=0}^n \mathcal{L}_k^{(-1/2)}(0) \left[\frac{\lambda-1}{\lambda+1} \frac{\lambda \cos^2(\phi/2) - \sin^2(\phi/2)}{\lambda \cos^2(\phi/2) + \sin^2(\phi/2)} \right]^{n-k} \times \mathcal{L}_{n-k}^{(-1/2)} \left[-\frac{2\lambda^2 p^2}{(\lambda^2 - 1)[\lambda^2 \cos^4(\phi/2) - \sin^4(\phi/2)]} \right] \exp \left[-\frac{\lambda p^2}{(\lambda+1)[\lambda \cos^2(\phi/2) + \sin^2(\phi/2)]} \right]. \quad (35)$$

Figures 1(a)-(d) show the three-dimensional plots of $P_n(p, q; \lambda, \phi)$ versus n and ϕ , for $\lambda = 21, 201, 1/21, 1/201$, respectively. The plane $\phi = 0$ in figure 1(a) corresponds to the oscillations pointed out in [1], which depend strongly on ϕ , showing a periodicity of π . Now, for $\lambda = 201$ [figure 1(b)] we observe the occurrence of rich structures, although the beats pointed out in [4] can not be perceived. In fact, they are revealed in figures 2 and 3. Figures 2(a)-(f) show the plots of P_n versus n for $\phi = 85^\circ, \dots, 90^\circ$ and $\lambda = 201$, where the beat structure becomes evident; however, it disappears at angles close to 90° . Following the arguments presented in [4] and corroborated by Mandal [7], this beat structure is a consequence of the quantum interference in phase space. Figures 3(a)-(f) show the plots of C_n versus n for the same parameters used in figures 2(a)-(f), where the beat structure is present again. This fact connects the *correlations* and *interference* effects in phase space, and provides further insights to the phenomenon. Moreover, we observe a similar kind of plots for $\lambda = 1/21$ and $\lambda = 1/201$, figures 1(c) and 1(d), respectively, where now they are shifted by $\pi/2$.

V. SUMMARY AND CONCLUSIONS

We have considered the Husimi function $P(p, q; \lambda, \phi)$ with emphasis on the marginal and correlation distribution functions, showing that all three satisfy the pseudo-diffusion equations if one considers that the squeezing parameter λ plays the role of a time. The solution, obtained from the Fourier transform method, permits calculating $P(p, q; \lambda, \phi)$, given an ‘initial’ Husimi function $P(p, q)$ with a kernel $K(\xi, \eta; \lambda, \phi)$ responsible by the propagation of squeezing. The decomposition of the kernel in three factors, equation (23), permits writing the CDF as a sum of two terms, having different interpretations: the first term, equation (26), is the propagation of ‘initial’ correlations contained in $P(p, q)$; whereas the second term, equation (27), is responsible for introducing additional correlations into the uncorrelated ‘initial’ product of the MDFs $Q(q)R(p)$.

Finally we remind that the formal procedure employed throughout this paper is advantageous if compared with the direct and lengthy calculation exposed in appendix A. In the specific case of the number state, the decomposition of the CDF into two terms should permit to investigate more thoroughly the origin of beats. Although having attained a formal expression for both (see appendix B), the numerical calculation presents difficulties due to its complexity.

Multimode-squeezed-states representation can also be considered within the present formalism. In particular, M. Salvadoray *et al.* [15] studied the two-mode-squeezed-state photon distribution. Again they verified the presence of beats. In this case, the correlation distribution function plays a crucial role in the understanding of this effect.

ACKNOWLEDGMENTS

MAM acknowledges financial support from FAPESP, São Paulo, project number 97/14551-4. SSM acknowledges financial support from CNPq, Brasil. This work has also been partially supported by Convênio FINEP/PRONEX Grant number 41/96/0935/00.

APPENDIX A: MDFs FOR THE NUMBER STATES BY DIRECT INTEGRATION

The usual procedure to calculate the MDFs for the number states consists in the integration of equation (30) with respect the variables p or q , respectively,

$$Q_n(q; \lambda, \phi) = \int_{-\infty}^{\infty} \frac{dp}{\sqrt{2\pi}} P_n(p, q; \lambda, \phi), \quad (\text{A1})$$

$$R_n(p; \lambda, \phi) = \int_{-\infty}^{\infty} \frac{dq}{\sqrt{2\pi}} P_n(p, q; \lambda, \phi). \quad (\text{A2})$$

We calculate (A1) by direct integration and present some properties inherent to the marginal distributions.

Consider initially the integral representation of the Hermite polynomial [16]

$$\mathcal{H}_n(z) = (-2i)^n e^{z^2} \int_{-\infty}^{\infty} \frac{du}{\sqrt{\pi}} u^n e^{-u^2 + 2iuz},$$

which permits to write (30) in a more convenient form:

$$\begin{aligned} P_n(p, q; \lambda, \phi) &= \frac{2\sqrt{\lambda}}{\lambda+1} \frac{(-2)^n}{n!} \left(\frac{\lambda-1}{\lambda+1} \right)^n \exp \left\{ - \left[\frac{\sin^2(\phi/2) - \lambda \cos^2(\phi/2)}{\lambda-1} \right] q^2 \right\} \\ &\times \exp \left\{ - \left[\frac{\cos^2(\phi/2) - \lambda \sin^2(\phi/2)}{\lambda-1} \right] p^2 + \left(\frac{\lambda+1}{\lambda-1} \sin \phi \right) pq \right\} \\ &\times \int_{-\infty}^{\infty} \frac{dudv}{\pi} (uv)^n \exp \left\{ -(u^2 + v^2) + \frac{2}{\sqrt{\lambda^2-1}} [(u-v) \sin(\phi/2) + i\lambda(u+v) \cos(\phi/2)] q \right\} \\ &\times \exp \left\{ - \frac{2}{\sqrt{\lambda^2-1}} [(u-v) \cos(\phi/2) - i\lambda(u+v) \sin(\phi/2)] p \right\}. \end{aligned} \quad (\text{A3})$$

Substituting (A3) into (A1) and integrating with respect to p , we obtain

$$\begin{aligned} Q_n(q; \lambda, \phi) &= \sqrt{2\alpha^2} \frac{(-2)^n}{n!} \left(\frac{\alpha}{\beta} \right)^n e^{-(\alpha q)^2} \int_{-\infty}^{\infty} \frac{dy}{\sqrt{\pi}} y^n e^{-(y-i\alpha q)^2} \\ &\times \int_{-\infty}^{\infty} \frac{dx}{\sqrt{\pi}} x^n \exp \left\{ - \left[x - i \left(\beta q + i \frac{\beta}{2\alpha} \left(1 - \frac{\alpha^2}{\beta^2} \right) y \right) \right]^2 \right\} \end{aligned} \quad (\text{A4})$$

for values of the squeeze parameter in the intervals $1 < \lambda < \cot^2(\phi/2)$ or $\cot^2(\phi/2) < \lambda < 1$, where

$$\alpha = \sqrt{\frac{\lambda}{(\lambda+1) [\cos^2(\phi/2) + \lambda \sin^2(\phi/2)]}} \quad \text{and} \quad \beta = \sqrt{\frac{\lambda}{(\lambda-1) [\cos^2(\phi/2) - \lambda \sin^2(\phi/2)]}}.$$

Integration over the variable x leads to [12, §3.462-4]

$$Q_n(q; \lambda, \phi) = \sqrt{2\alpha^2} \frac{(-i)^n}{n!} \left(\frac{\alpha}{\beta} \right)^n e^{-(\alpha q)^2} \int_{-\infty}^{\infty} \frac{dy}{\sqrt{\pi}} y^n \mathcal{H}_n \left[\beta q + i \frac{\beta}{2\alpha} \left(1 - \frac{\alpha^2}{\beta^2} \right) y \right] e^{-(y-i\alpha q)^2}. \quad (\text{A5})$$

Now, using relation [14]

$$\mathcal{H}_n(z+w) = \sum_{k=0}^n \mathcal{L}_{n-k}^{(k)}(0) (2w)^{n-k} \mathcal{H}_k(z)$$

for the Hermite polynomial present in (A5),

$$\mathcal{H}_n \left[\beta q + i \frac{\beta}{2\alpha} \left(1 - \frac{\alpha^2}{\beta^2} \right) y \right] = \sum_{k=0}^n \mathcal{L}_{n-k}^{(k)}(0) \left[i \frac{\beta}{\alpha} \left(1 - \frac{\alpha^2}{\beta^2} \right) y \right]^{n-k} \mathcal{H}_k(\beta q),$$

we get

$$Q_n(q; \lambda, \phi) = \frac{\sqrt{2\alpha^2}}{2^n n!} \left(\frac{\alpha}{\beta} \right)^n e^{-(\alpha q)^2} \sum_{k=0}^n \mathcal{L}_{n-k}^{(k)}(0) \left[\frac{\beta}{2\alpha} \left(\frac{\alpha^2}{\beta^2} - 1 \right) \right]^{n-k} \mathcal{H}_k(\beta q) \mathcal{H}_{2n-k}(\alpha q). \quad (\text{A6})$$

Note that (A6) can be written in a compact form and equivalent to equation (34), *i.e.*, in terms of the associated Laguerre polynomial. For this purpose it is necessary to verify the equality

$$\frac{1}{2^{2n} n!} \sum_{s=0}^n \mathcal{L}_{n-s}^{(s)}(0) \left(\frac{2\alpha}{\beta} \right)^s \left(\frac{\alpha^2}{\beta^2} - 1 \right)^{n-s} \mathcal{H}_s(\beta q) \mathcal{H}_{2n-s}(\alpha q) = \sum_{k=0}^n c_{nk}(\alpha, \beta) \mathcal{L}_k^{(-1/2)} [2(\alpha\beta q)^2] \quad (\text{A7})$$

and to determine the coefficients $c_{nk}(\alpha, \beta)$.

The relation established by Bailey [17] for the product of Hermite polynomials,

$$\begin{aligned} \mathcal{H}_m(ax) \mathcal{H}_l(bx) &= \sum_{j=0}^{\lfloor \frac{m+l}{2} \rfloor} (-1)^j \frac{m!}{j!(m-2j)!} \frac{a^{m-2j} b^{2j-l}}{(\sqrt{a^2+b^2})^{m-l}} {}_2F_1 \left(m+1, -l; m-2j+1; \frac{a^2}{a^2+b^2} \right) \\ &\times \mathcal{H}_{m+l-2j}(\sqrt{a^2+b^2} x) \end{aligned}$$

where ${}_2F_1(a_1, a_2; a_3; z)$ is the hypergeometric function, permits to verify the equality (A7) through

$$\begin{aligned} \mathcal{H}_s(\beta q) \mathcal{H}_{2n-s}(\alpha q) &= \frac{(-1)^n (2n-s)!}{2^{n-s} (\alpha\beta)^{2n-s}} \sum_{k=0}^n \frac{k!}{(2k-s)!(n-k)!} (2\alpha)^{2k} \beta^{2(n-k)} \\ &\times {}_2F_1 \left(2n-s+1, -s; 2k-s+1; \frac{1}{2\beta^2} \right) \mathcal{L}_k^{(-1/2)} [2(\alpha\beta q)^2], \end{aligned} \quad (\text{A8})$$

and to determine the coefficients $c_{nk}(\alpha, \beta)$,

$$\begin{aligned} c_{nk}(\alpha, \beta) &= \frac{(-1)^n}{2^{2n}} \frac{k!}{(n-k)!} \frac{(2\alpha)^{2k} \beta^{2(n-k)}}{(2\alpha^2 \beta^2)^n} \sum_{s=0}^n \frac{(2n-s)!}{s!(n-s)!(2k-s)!} (2\alpha)^{2s} \left(\frac{\alpha^2}{\beta^2} - 1 \right)^{n-s} \\ &\times {}_2F_1 \left(2n-s+1, -s; 2k-s+1; \frac{1}{2\beta^2} \right). \end{aligned} \quad (\text{A9})$$

In fact, the sum present in (A9) can be performed since we used the following relation for the hypergeometric function [14]:

$${}_2F_1 \left(2n-s+1, -s; 2k-s+1; \frac{1}{2\beta^2} \right) = \frac{s!(2k-s)!}{(2n-s)!} \sum_{l=0}^s (-1)^l \frac{(2n-s+l)!}{(s-l)!(2k-s+l)!} \frac{(2\beta^2)^{-l}}{l!}. \quad (\text{A10})$$

Then, substituting (A10) into (A9) and calculating the sums, we obtain a simple expression for the coefficients

$$c_{nk}(\alpha, \beta) = (-1)^k \frac{[2(n-k)-1]!!}{[2(n-k)]!!} \left(\frac{\alpha}{\beta} \right)^{2k} = (-1)^k \mathcal{L}_{n-k}^{(-1/2)}(0) \left(\frac{\alpha}{\beta} \right)^{2k}. \quad (\text{A11})$$

So, the marginal distribution (A6) can be expressed such as equation (34). Adopting analogous procedure for equation (A2) we obtain (35), however the values of squeeze parameter are restricted in the intervals $\tan^2(\phi/2) > \lambda > 1$ or $\tan^2(\phi/2) < \lambda < 1$.

In addition to the symmetry relations established in section III [see equation (22)], the MDFs exhibit the following properties:

$$(i) \lim_{\lambda \rightarrow \infty} Q_n(q; \lambda, 0) = |\Psi_n(q)|^2 \quad \text{and} \quad \lim_{\lambda \rightarrow 0} R_n(p; \lambda, 0) = |\Phi_n(p)|^2, \quad (\text{A12})$$

$$(ii) \sum_{n=0}^{\infty} Q_n(q; \lambda, \phi) = \sum_{n=0}^{\infty} R_n(p; \lambda, \phi) = \sum_{n=0}^{\infty} \mathcal{L}_n^{(-1/2)}(0) \rightarrow \infty. \quad (\text{A13})$$

The first property does not characterize the Husimi function as a probability distribution but only emphasizes the character of quasiprobability distribution: limit values of λ recover the squared moduli of wavefunctions [10,18]. With respect to the second property, it is a direct consequence of the scalar product for the squeezed states, *i.e.*, $\sum_{n=0}^{\infty} P_n(p, q; \lambda, \phi) = 1$. In fact, the divergence is a consequence of the integration step of this relation over the variables p or q . Now, considering the normalization of equation (30), $\int_{-\infty}^{\infty} \frac{dpdq}{2\pi} P_n(p, q; \lambda, \phi) = 1$, we obtain the third property for MPDFs:

$$(iii) \int_{-\infty}^{\infty} \frac{dq}{\sqrt{2\pi}} Q_n(q; \lambda, \phi) = 1 \quad \text{and} \quad \int_{-\infty}^{\infty} \frac{dp}{\sqrt{2\pi}} R_n(p; \lambda, \phi) = 1. \quad (\text{A14})$$

APPENDIX B: THE COMPONENTS OF THE CORRELATION DISTRIBUTION FUNCTION

The components of the correlation distribution function $C(p, q; \lambda, \phi)$ can also be written as

$$C^{(1)}(p, q; \lambda, \phi) = P(p, q; \lambda, \phi) - C^{(3)}(p, q; \lambda, \phi), \quad (\text{B1})$$

$$C^{(2)}(p, q; \lambda, \phi) = C^{(3)}(p, q; \lambda, \phi) - Q(q; \lambda, \phi)R(p; \lambda, \phi), \quad (\text{B2})$$

with

$$C^{(3)}(p, q; \lambda, \phi) = \int_{-\infty}^{\infty} \frac{d\xi d\eta}{2\pi} e^{i(\eta p - \xi q)} K(\xi, \eta; \lambda, \phi) \tilde{Q}(\xi; 1) \tilde{R}(\eta; 1). \quad (\text{B3})$$

Present in both equations (B1) and (B2), the term $C^{(3)}$ is responsible for introducing correlations into the marginal distributions product through the relation

$$C^{(3)}(p, q; \lambda, \phi) = \exp \left[\left(\frac{\lambda^2 - 1}{4\lambda} \sin \phi \right) \frac{\partial^2}{\partial p \partial q} \right] Q(q; \lambda, \phi) R(p; \lambda, \phi), \quad (\text{B4})$$

where we used the result:

$$e^{i(\eta p - \xi q)} K(\xi, \eta; \lambda, \phi) = \exp \left[\left(\frac{\lambda^2 - 1}{4\lambda} \sin \phi \right) \frac{\partial^2}{\partial p \partial q} \right] e^{i(\eta p - \xi q)} k_Q(\xi; \lambda, \phi) k_R(\eta; \lambda, \phi).$$

For $\phi = 0$, equations (B1)-(B3) simplify to

$$\begin{aligned} C^{(1)}(p, q; \lambda, 0) &= C(p, q; \lambda, 0), \\ C^{(2)}(p, q; \lambda, 0) &= 0, \\ C^{(3)}(p, q; \lambda, 0) &= Q(q; \lambda, 0)R(p; \lambda, 0). \end{aligned}$$

In this appendix we calculate the component $C^{(3)}$ for the number states and, consequently, the components $C^{(1)}$ and $C^{(2)}$ can be totally determined.

In order to simplify the calculations, let us initially consider the marginal and correlation distribution functions expressed in terms of the parameters α_1 , α_2 and α_3 :

$$Q_n(q; \lambda, \phi) = \sqrt{2\alpha_1^2} e^{-(\alpha_1 q)^2} \sum_{k=0}^n \mathcal{L}_{n-k}^{(k)}(0) (-2\alpha_1^2)^k \mathcal{L}_k^{(-1/2)}[(\alpha_1 q)^2], \quad (\text{B5})$$

$$R_n(p; \lambda, \phi) = \sqrt{2\alpha_2^2} e^{-(\alpha_2 p)^2} \sum_{k=0}^n \mathcal{L}_{n-k}^{(k)}(0) (-2\alpha_2^2)^k \mathcal{L}_k^{(-1/2)}[(\alpha_2 p)^2], \quad (\text{B6})$$

$$C_n^{(3)}(p, q; \lambda, \phi) = \sum_{l=0}^{\infty} \frac{\alpha_3^l}{l!} \frac{\partial^{2l}}{\partial p^l \partial q^l} Q_n(q; \lambda, \phi) R_n(p; \lambda, \phi), \quad (\text{B7})$$

with

$$\begin{aligned}\alpha_1 &= \sqrt{\frac{\lambda}{(\lambda+1) [\cos^2(\phi/2) + \lambda \sin^2(\phi/2)]}} , \\ \alpha_2 &= \sqrt{\frac{\lambda}{(\lambda+1) [\lambda \cos^2(\phi/2) + \sin^2(\phi/2)]}} , \\ \alpha_3 &= \frac{\lambda^2 - 1}{4\lambda} \sin \phi ,\end{aligned}$$

and $\alpha_1^2 + \alpha_2^2 = 2\alpha_1^2\alpha_2^2$. Expressions (B5) and (B6) are the alternative way of writing the marginal distributions (34) and (35), respectively, since we used the properties of the associated Laguerre polynomials. Moreover, using the relation

$$\frac{\partial^l}{\partial x^l} \left\{ e^{-(ax)^2} \mathcal{L}_k^{(-1/2)}[(ax)^2] \right\} = \frac{(-1)^k}{2^{2k} k!} (-a)^l e^{-(ax)^2} \mathcal{H}_{l+2k}(ax) ,$$

those expressions permit to calculate the component $C_n^{(3)}$, given the result:

$$\begin{aligned}C_n^{(3)}(p, q; \lambda, \phi) &= 2\alpha_1\alpha_2 e^{-[(\alpha_1 q)^2 + (\alpha_2 p)^2]} \sum_{k=0}^n \mathcal{L}_{n-k}^{(k)}(0) \frac{\alpha_1^{2k}}{2^k k!} \sum_{m=0}^n \mathcal{L}_{n-m}^{(m)}(0) \frac{\alpha_2^{2m}}{2^m m!} \\ &\quad \times \sum_{l=0}^{\infty} \frac{(\alpha_1\alpha_2\alpha_3)^l}{l!} \mathcal{H}_{l+2k}(\alpha_1 q) \mathcal{H}_{l+2m}(\alpha_2 p) .\end{aligned}\tag{B8}$$

The infinity sum present in (B8) can be done by using the formula [19, §5.12.2.1]

$$\begin{aligned}\sum_{k=0}^{\infty} \frac{t^k}{k!} \mathcal{H}_{k+m}(x) \mathcal{H}_{k+n}(y) &= \frac{1}{\sqrt{(1-4t^2)^{m+n+1}}} \exp \left[\frac{4txy - 4t^2(x^2 + y^2)}{1-4t^2} \right] \sum_{r=0}^{\{m,n\}} r! \mathcal{L}_{m-r}^{(r)}(0) \mathcal{L}_{n-r}^{(r)}(0) (4t)^r \\ &\quad \times \mathcal{H}_{m-r} \left(\frac{x-2ty}{\sqrt{1-4t^2}} \right) \mathcal{H}_{n-r} \left(\frac{y-2tx}{\sqrt{1-4t^2}} \right) \quad |t| < 1/2 ,\end{aligned}$$

where $\{m, n\}$ stands for the minor of m and n , which leads to

$$\begin{aligned}C_n^{(3)}(p, q; \lambda, \phi) &= \frac{2\alpha_1\alpha_2}{\sqrt{1-(2\alpha_1\alpha_2\alpha_3)^2}} \exp \left[-\frac{\alpha_1^2 q^2 - 4\alpha_1^2\alpha_2^2\alpha_3 pq + \alpha_2^2 p^2}{1-(2\alpha_1\alpha_2\alpha_3)^2} \right] \sum_{k=0}^n \frac{\mathcal{L}_{n-k}^{(k)}(0)}{2^k k!} \left[\frac{\alpha_1^2}{1-(2\alpha_1\alpha_2\alpha_3)^2} \right]^k \\ &\quad \times \sum_{m=0}^n \frac{\mathcal{L}_{n-m}^{(m)}(0)}{2^m m!} \left[\frac{\alpha_2^2}{1-(2\alpha_1\alpha_2\alpha_3)^2} \right]^m \sum_{r=0}^{\{2k, 2m\}} r! \mathcal{L}_{2k-r}^{(r)}(0) \mathcal{L}_{2m-r}^{(r)}(0) (4\alpha_1\alpha_2\alpha_3)^r \\ &\quad \times \mathcal{H}_{2k-r} \left(\frac{\alpha_1 q - 2\alpha_1\alpha_2^2\alpha_3 p}{\sqrt{1-(2\alpha_1\alpha_2\alpha_3)^2}} \right) \mathcal{H}_{2m-r} \left(\frac{\alpha_2 p - 2\alpha_1^2\alpha_2\alpha_3 q}{\sqrt{1-(2\alpha_1\alpha_2\alpha_3)^2}} \right) .\end{aligned}\tag{B9}$$

- [1] Schleich W and Wheeler J A 1987 *J. Opt. Soc. Am. B* **4** 1715
Schleich W and Wheeler J A 1987 *Nature* **326** 574
Schleich W, Walls D F and Wheeler J A 1988 *Phys. Rev. A* **38** 1177
- [2] Vourdas A and Weiner R M 1987 *Phys. Rev. A* **36** 5866
Dodonov V V, Klimov A B and Man'ko V I 1989 *Phys. Lett. A* **134** 211
- [3] Gagen M J 1995 *Phys. Rev. A* **51** 2715
- [4] Dutta B, Mukunda N, Simon R and Subramanian A 1993 *J. Opt. Soc. Am. B* **10** 253

- [5] Chountasis S and Vourdas A 1998 *Phys. Rev. A* **58** 848
Chountasis S and Vourdas A 1998 *Phys. Rev. A* **58** 1794
- [6] Daboul J, Marchiolli M A and Mizrahi S S 1995 *J. Phys. A: Math. Gen.* **28** 4623
Mizrahi S S and Marchiolli M A 1996 *IV Wigner Symposium: Proc. Symp. (Mexico, 1995)* ed N M Atakishiyev, T H Seligman and K B Wolf (Singapore: World Scientific) p 158
Daboul J 1996 *Phys. Lett. A* **212** 1
- [7] Mandal S 1998 *Phys. Rev. A* **58** 752 and references therein
- [8] Mizrahi S S and Daboul J 1992 *Physica A* **189** 635
Daboul J and Mizrahi S S 1994 *Journal of Group Theory in Physics* **2** 161
- [9] Mizrahi S S and Marchiolli M A 1993 *Physica A* **199** 96
- [10] Mizrahi S S 1984 *Physica A* **127** 241
- [11] Louisell W H 1990 *Quantum Statistical Properties of Radiation* (New York: Wiley)
Mandel L and Wolf E 1995 *Optical Coherence and Quantum Optics* (New York: Cambridge Univ. Press)
Scully M O and Zubairy M S 1997 *Quantum Optics* (New York: Cambridge Univ. Press)
- [12] Gradshteyn I S and Ryzhik I M 1980 *Table of Integrals, Series and Products* (New York: Academic Press)
- [13] Yuen H P 1976 *Phys. Rev. A* **13** 2226
- [14] Magnus W, Oberhettinger F and Soni R P 1966 *Formulas and Theorems for the Special Functions of Mathematical Physics* (New York: Springer Verlag)
- [15] Salvadoray M, Kumar M S and Simon R 1994 *Phys. Rev. A* **49** 4957 and references therein
- [16] Lebedev N N 1972 *Special Functions and their Applications* (New York: Dover)
Arfken G 1985 *Mathematical Methods for Physicists* (San Diego: Academic Press)
- [17] Bailey W N 1948 *J. London Math. Soc.* **23** 291
- [18] Mizrahi S S 1988 *Physica A* **150** 541
Mizrahi S S and Galetti D 1988 *Physica A* **153** 567
- [19] Prudnikov A P, Brychkov Y A and Marichev O I 1986 *Integrals and Series: Special Functions* (New York: Gordon and Breach Science Publishers)

FIG. 1. Plots of $P_n(p, q; \lambda, \phi)$ vs. n and ϕ with $p^2 + q^2 = 98$. Figures 1(a)-(d) correspond to $\lambda = 21, 201, 1/21$ and $1/201$, respectively.

FIG. 2. Plots of $P_n(p, q; \lambda, \phi)$ vs. n with $\phi = 85^\circ, \dots, 90^\circ$ and $\lambda = 201$. The phase-space variables are transformed into $q = 7\sqrt{2} \cos \theta$ and $p = 7\sqrt{2} \sin \theta$, which leads to $q_r = 7\sqrt{2} \cos(\theta - \phi/2)$ and $p_r = 7\sqrt{2} \sin(\theta - \phi/2)$. For mapping the parameters used by Dutta *et al.*, we fixed $\theta = 3\phi/2$.

FIG. 3. Plots of $C_n(p, q; \lambda, \phi)$ vs. n for the same parameters set used in figure 2, where we see the presence of beats again. This fact corroborates the phase-space interference concept since *correlations* and *interference* effects are closely connected.